

QUESTIONS & ANSWERS

Kill your exam at first Attempt



NCAC-I Dumps
NCAC-I Braindumps
NCAC-I Real Questions
NCAC-I Practice Test
NCAC-I dumps free



NCC-AP

NCAC-I

National Certified Addiction Counselor, Level I

<https://killexams.com/pass4sure/exam-detail/NCAC-I>



Question: 194

What is a common medication used for the treatment of nicotine use disorder that helps reduce cravings and withdrawal symptoms by providing nicotine replacement?

- A. Varenicline
- B. Naltrexone
- C. Acamprosate
- D. Nicotine gum

Answer: D

Question: 195

Which of the following is a technique used in relapse prevention for substance use disorders that involves developing a plan for managing high-risk situations?

- A. Assertiveness training
- B. Cue exposure therapy
- C. Contingency management
- D. Coping skills training

Answer: D

Question: 196

What is a common symptom of cannabis withdrawal?

- A. Mydriasis
- B. Hypotension
- C. Hyperactivity
- D. Irritability

Answer: D

Question: 197

Which of the following is a common symptom of sedative-hypnotic use disorder?

- A. Mydriasis
- B. Hypotension
- C. Hyperactivity
- D. Sedation

Answer: D

Question: 198

What is a common medication used for the treatment of stimulant use disorder that helps reduce cravings and withdrawal symptoms by affecting norepinephrine levels?

- A. Methadone
- B. Naloxone
- C. Buprenorphine
- D. Atomoxetine

Answer: D

Question: 199

Which of the following is a technique used in relapse prevention for substance use disorders that involves developing a relapse prevention plan and practicing new coping skills?

- A. Assertiveness training
- B. Cue exposure therapy
- C. Contingency management
- D. Coping skills training

Answer: D

Question: 200

What is a common symptom of caffeine withdrawal?

- A. Mydriasis
- B. Hypotension
- C. Fatigue
- D. Hyperactivity

Answer: C

Question: 201

What is a common medication used for the treatment of sedative-hypnotic use disorder that helps reduce cravings and withdrawal symptoms by modulating glutamate activity?

- A. Methadone
- B. Naloxone
- C. Buprenorphine
- D. Gabapentin

Answer: D

Question: 202

Which of the following is a technique used in motivational interviewing for substance use disorders that involves exploring the client's confidence in making changes?

- A. Reflective listening
- B. Decisional balance
- C. Directive advice
- D. Scaling questions

Answer: D

Question: 203

What is a common symptom of hallucinogen withdrawal?

- A. Mydriasis
- B. Hypertension
- C. Hyperactivity
- D. Depressed mood

Answer: D



For More exams visit <https://killexams.com/vendors-exam-list>



THESE ARE SAMPLE QUESTIONS

**Full Version Contains
UP-TO-DATE
Actual Questions and Verified
Answers**